



New Horizons Journal of Basic and Applied Sciences

Monitoring aflatoxins contamination in maize using chemical treatments

Marwa A. Mahmoud, Abdallah M. A. Hassane* and Nageh F. Abo-Dahab

Botany and Microbiology Department, Faculty of Science, Al-Azhar University, Assiut 71524, Egypt

Received: 04, 05, 2026; Accepted: 23, 05, 2026; Published: 19, 07, 2026

© 2026 The Author(s). Published by Science Park Publisher. This is an open access article under the CC BY 4.0 license (<https://creativecommons.org/licenses/by/4.0/>)

Abstract

Aflatoxins (AFs) contamination, which is primarily driven by *Aspergillus flavus* and *A. parasiticus*, represents a global threat to food safety and the agricultural economy, with maize (*Zea mays* L.) being one of the most susceptible staple crops. These secondary metabolites are potent carcinogens, linked to hepatocellular carcinoma and immunosuppression among others. Contamination is heavily influenced by environmental stressors, particularly drought and high temperatures, which are increasingly prevalent due to global climate change. While pre-harvest and post-harvest management strategies such as biocontrol and secure storage have shown efficacy, they are often insufficient to prevent AFs contamination. Oxidative stress is closely associated with aflatoxin biosynthesis, where the fungus produces toxins as a defensive mechanism against reactive oxygen species in its environment. Consequently, the application of chemical agents has emerged as a promising strategy for mitigation. Natural and synthetic antioxidants can suppress *Aspergillus* growth and reduce toxin accumulation in stored grains. Emerging techniques also highlight the use of antioxidant-rich phenolic compounds as redox mediators to enhance the enzymatic disruption of AF biosynthetic pathway. This review concludes that integrating antioxidant-based interventions into holistic farm-to-market management systems is vital for improving maize safety and protecting vulnerable populations.

Keywords: Aflatoxins; Maize; *Aspergillus*; Antioxidants; Food safety

1. Introduction

Mycotoxins are toxic secondary metabolites that are synthesized by some fungal genera, like *Fusarium*, *Aspergillus*, and *Penicillium*, which contaminants estimated between 60 and 80% of the world's food crops [1]. The most common mycotoxins are aflatoxins (carcinogens found in nuts and dairy products), ochratoxins (common in coffee and grains), zearalenone (an estrogenic toxin in cereals), fumonisins (in maize), trichothecenes (causing aleukia), and patulin (in apple products) [2]. These compounds are dangerous to human and animal health because they are carcinogenic, mutagenic, teratogenic, and immunosuppressive [3]. Environmental factors like moisture and high temperatures are the main factors that promote fungal growth and toxin production that are common during storage

processes, transport, and pre-harvest [4]. Since mycotoxins are stable at high temperatures and toxic even at low concentrations, a lot of countries have set regulatory limits on mycotoxins in food and feed products, usually between 2 and 20 part per billion (ppb) [5].

Maize (*Zea mays* L.) Family Poaceae is one of the most economically important cereal crops across the globe, a major source of food, nutrition security, and income to commercial farmers [6]. Over 197 million hectares of land worldwide are under maize farming and yield about 1.13 billion tons [7]. It enables it to find wide application in industries in human consumption and bio energy, like the production of bioethanol [8]. Maize is very susceptible to infection by toxigenic fungi belonging to the genera *Aspergillus* and *Fusarium* and contamination by their toxigenic mycotoxins during its storage, harvesting, and growth [6]. Climate change is linked

Review Article

to this vulnerability since the conditions of the environment, such as heat and unpredictable precipitation occurrence during periods of the crop lifecycle, such as grain filling and silking, can decrease the yields and increase the toxin level [9]. To reduce the growth of molds, such as *Aspergillus* and *Fusarium* as major producers of mycotoxins that spoil grains by producing mycotoxins, it is important to dry the grains [10]. Since maize is an important component of the global food and feed chain, ensuring its quality and protection against these biological and environmental hazards is a major concern for global food security [7]. Despite the vast number of published articles and reviews concerning mycotoxin contamination in diverse food and feedstuffs, the present review sheds light on aflatoxin contamination in maize and maize products with reference to chemical management of this contaminant, which has not been addressed previously with this particularity.

2. Mycoflora of maize

The mycoflora of maize is famous for a rich number of

fungal species involved in contaminating the crop both at post-harvest and pre-harvest levels [11]. According to Nesci et al. [12], *Aspergillus*, *Fusarium*, and *Penicillium* genera were found to be the most maize-related fungal genera. The frequency of fungi in maize kernels indicated that *Aspergillus flavus* is the most common (52%), then *Fusarium* spp. (25%) and *Penicillium* spp. (21%) [13]. Some typical isolates of infected maize grains are *Aspergillus flavus*, *A. parasiticus*, *A. niger*, and *Rhizopus nigricans* [14]. These organisms are the main sources of spoilage contamination, cause loss of the grain quality, and deteriorate proteins [15]. This mycoflora is responsible for the toxic synthesis and carcinogenic secondary metabolites, such as AFs and fumonisins, dangerous to health and humans' life [2]. Saleh et al. [14] recorded that *A. parasiticus* (32.5%) and *A. flavus* (47.5%) were the dominant fungi in corn with low levels of AFs (AFB₁, AFB₂, AFG₁ and AFG₂) all under international safety limits.




Figure 1. Chemical structure of aflatoxins adopted from Hassane [24].

Review Article

3. Aflatoxins

Aflatoxins are very toxic and carcinogenic secondary metabolites that are produced by filamentous fungi of the genus *Aspergillus*, like *A. flavus* and *A. parasiticus*, which are contaminants of crops [16]. Environmental factors affect their production and grow under warm and humid conditions with temperatures best between 25°C and 35°C and high moisture content [17]. These toxins are classified into four primary types—AFB₁, AFG₁, AFB₂, and AFG₂—named for the green (G) or blue (B) fluorescence they emit under ultraviolet light, alongside their hydroxylated metabolites AFM₁ and AFM₂, which are commonly detected in animal products such as milk [18] (Figure 1). Among these congeners, AFB₁ is identified as the most potent and prevalent toxicant, because of its direct link to liver cancer and its ability to cause acute aflatoxicosis, with the relative toxicity ranking generally established as B₁ > G₁ > B₂ > G₂ [5, 19]. Due to their severe health impacts, including mutagenic, teratogenic, and immunosuppressive effects, AFs are classified as Group 1 human carcinogens by the International Agency for Research on Cancer (IARC) [20]. AFB₁ causes acute/chronic aflatoxicosis, hepatocellular

carcinoma, immunosuppression, and stunted growth in children. In milk, AFM₁ is concern to the infants [21].

The synthesis of AFs is an enzymatic process that requires the expression of genes such as *aflP*, *aflE*, and *aflQ*, which transform precursors such as acetyl-CoA into AFB₁, AFG₁, AFB₂, and AFG₂. AFB₁ is metabolized to AFB₁-8,9-epoxide, leading to DNA damage and liver cancer [22]. Exposure to humans and animals by inhalation, ingestion, or skin contact, and linked to clinical effects including lethal acute aflatoxicosis, liver cancer, and growth retardation in children [23]. Aflatoxins occur in a variety of agricultural products, like maize, wheat, rice, peanuts, and spices, and cause economic losses and a high risk to world food security [17].

4. Aflatoxin biosynthesis

Aflatoxin biosynthesis in *Aspergillus* was governed by a cluster which includes 30 genes, where its activation is regulated mainly by *aflR* and *aflS* genes, and at least 27 enzymatic processes have been implicated [25, 26]. Table (1) and Figure (2) illustrate a summary of the consequences in the AFB₁ biosynthesis pathway.




Figure 2. Aflatoxin B1 pathway. Numbers and abbreviations refer to enzymes stated in table 1.

Review Article

Table 1. Aflatoxin B₁ pathway cluster genes and enzymes involved in biosynthesis.

No.	Compound/Substrate	Abbreviation	Enzyme	Gene	Action
1	Acetyl-CoA & Malonyl-CoA	-	Fatty acid synthase	<i>aflA</i> & <i>aflB</i>	Condensation
2	Hexanoate	-	Polyketide synthase	<i>aflC</i>	Chain elongation
3	Polyketide	-	Polyketide synthase	<i>aflC</i>	Cyclization
4	Norsolorinic acid	NOR	Reductase	<i>aflD</i>	Reduction
5	Averantin	AVN	P450 monooxygenase	<i>aflG</i>	Oxidation
6	5'-Hydroxyaverantin	HAVN	Alcohol dehydrogenase	<i>aflH</i>	Oxidation
7	5'-Oxoaverantin	OAVN	Synthase	<i>aflK</i>	Bisfuran ring closure
8	Averufin	AVF	Oxidase	<i>aflV</i> , <i>aflI</i> , & <i>aflW</i>	Oxidation
9	Versiconal hemiacetal acetate	VHA	Esterase	<i>aflJ</i>	Cyclodehydration
10	Versiconal	VAL	VERB synthase	<i>aflK</i>	Bisfuran ring closure
11	Versicolorin B	VERB	Desaturase	<i>aflL</i>	Oxidation
12	Versicolorin A	VERA	Dehydrogenase/ketoreductase Monooxygenase	<i>aflN</i> , <i>aflM</i> , <i>aflY</i> , & <i>aflX</i>	Reduction
13	Demethylsterigmatocystin	DMST	O-Methyltransferase I / O-methyltransferase B	<i>aflO</i>	Methylation
14	Sterigmatocystin	ST	O-Methyltransferase A / O-methyltransferase II	<i>aflP</i>	Methylation
15	O-methylsterigmatocystin	OMST	Oxidoreductase/P450 monooxygenase	<i>aflQ</i> & <i>aflE</i>	Oxidation
16	Aflatoxin B ₁	AFB ₁	-	-	

5. Aflatoxigenic fungi

Aflatoxigenic fungi are primarily secondary metabolite producers belonging to the *Aspergillus* section *Flavi*, with *A. flavus* and *A. parasiticus* being the most prevalent species identified in global food and feed contamination [11]. The 24 *Aspergillus* species produce aflatoxins are divided into sections *Ochraceorosei*, *Flavi*, and *Nidulantes* [27]. Other producers are species like *A. parvisclerotigenus*, *A. nomius*, *A. bombycis*, *A. pseudotamarii*, and others of the section *Ochraceorosei* like *A. ostianus* [28, 29]. These fungi are common in air and soil, and they attack crops, such as maize, cotton, and peanuts during both post-harvest and pre-harvest stages [30]. Abiotic factors, like high temperatures (optimally 25–37°C) and high-water activity (more than 0.85 a_w), play an important role in their growth and production of toxins [31].

Toxigenic fungi, such as *A. flavus* and *A. parasiticus*, contaminated grains (corn, rice, and wheat). High-performance liquid chromatography (HPLC) and Thin-layer chromatography (TLC) were used to identify 21 of 22 isolates yielding AFs, including AFB₁, AFG₁, AFB₂, and AFG₂. The AFB₁ levels were between 0.02 and 875.03 ng/g, with the highest produced being AFG₂ [32, 33].

6. Environmental factors affecting aflatoxin production

The toxin production and growth of *Aspergillus* species are sensitive to biotic and abiotic environmental factors, with water activity (a_w) and temperature being the main factors that affect the production of toxins and fungal growth [5, 34]. Studies show that optimal temperature ranges for the

Review Article

production of aflatoxin are between 29°C and 33°C, and high-water activity of about 0.99 a_w [35]. Temperatures lower than 25°C or above 37°C are not conducive to biosynthesis, and production is slowed down at moisture below 0.85 a_w and between 0.70 and 0.75 a_w [5]. Agricultural substrates, such as moisture content, poultry feed is found to be the first limiting factor, with no biosynthesis at 11% moisture content, and growth increasing with moisture content to 17% moisture content [36]. Oxidative stress was identified as a critical prerequisite for aflatoxin production [37]. High temperature and drought, generated by climate change, are environmental stressors causing the buildup of reactive oxygen species (ROS) like hydrogen peroxide (H_2O_2) in the fungus [38]. The hypothesis is that the fungus induces biosynthesis of aflatoxin as a defense to counter these free radicals and reduce damage to the cell [3].

It has been suggested that aflatoxin production can have a functional role for the fungus, as ROS scavenging under oxidative stress and as a deterrent against frugivorous insects [11]. Such a relationship is supported by the fact that the expression of the aflatoxin gene cluster (*aflR* and *aflS*) in the environment is decimated by the antioxidant's introduction (caffeic acid, phenolic compounds, and selenium) and reduce toxin accumulation [37]. Also, gaseous composition (including CO_2 concentrations), pH (with the optimal range which is between 4 and 6), and the presence of nitrogen and carbon sources also control the toxigenicity of *Aspergillus* strains and metabolic activity [39].

7. Worldwide limits for aflatoxins in food

The control of AF in food and feed traces began in the late 1960s, and presently, about 120 nations have enacted specific regulatory limits to enable international trade and reduce health risks [40]. The multinational and national organizations set standards, and the Codex Alimentarius Commission, the United States Food and Drug Administration (US FDA), and the European Union (EU) are the main standards [41]. The EU has some limits in the world with 2 $\mu g/kg$ (ppb) for AFB_1 and 4 $\mu g/kg$ for total AFs in maize and peanuts that are eaten directly by humans [6]. The US FDA, like Brazil, China, and Ecuador, applies a higher level of 20 $\mu g/kg$ for total AFs in human food [42]. In the East African Community (EAC), there are regulatory thresholds that have been passed to ensure food safety, with the maximum permissible level of AFB_1 at 5

$\mu g/kg$ and total AFs at 10 $\mu g/kg$ [43]. The acceptable levels of Aflatoxin M1 in milk, with the EU setting the levels at 0.05 $\mu g/kg$, and the US setting the levels at 0.50 $\mu g/kg$ [42]. In these legal systems, the implementation of these rules in developing areas has been a main problem associated with food shortage, socioeconomic factors, and poor infrastructure [44].

8. Detection of aflatoxins

At the biological level, qualitative methods like UV fluorescence of special media, ammonia vapor test are cheap culture tools that allow separation of toxigenic and atoxigenic strains of fungi [45]. Identification of these strains is based on culture methods, such as green or blue fluorescence under UV light on coconut milk agar or detection of a color change when the vapors of ammonia are added [46], or modern molecular Polymerase Chain Reaction (PCR) assays [14]. Three types of analysis methods: immune-chemical, chromatographic, and spectroscopic used to detect and quantify AFs [47].

Several chromatographic and spectroscopic methods are utilized for detection and quantification of mycotoxins. Chromatographic techniques, including TLC, HPLC, and Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS), have been identified as reference standards because they are accurate [48]. TLC is a non-specific and low-cost screening method, but high-performance TLC (HPTLC) can enhance accuracy with specific mycotoxins such as AFs and ochratoxin A, but Gas chromatography (GC) is limited due to the low volatility of mycotoxins, though GC-MS is used for volatile mycotoxins like patulin. Liquid chromatography (LC), especially LC-MS/MS, is common in sensitive, selective, and has the capability of analyzing multiple mycotoxins at the same time without the need to be derivatized. The upgrade of sensitivity and separation efficiency is ultra-HPLC (UHPLC) and high-performance liquid chromatography (HPLC), which is combined with mass spectrometry detectors or fluorescence [49]. In the cutting-edge approach to detecting mycotoxins range and their metabolites in trace concentrations in complex food samples is LC-MS/MS [48]. Enzyme-Linked Immunosorbent Assay (ELISA) is the most used rapid immune-chemical method of high screening with high throughput, while immune-dipsticks and lateral flow devices (LFDs) are convenient and used during inspections [47]. Green methods of rapid, non-destructive, and real-time monitoring are explored using

Review Article

emerging and spectroscopic technologies, like Hyperspectral Imaging (HSI), Surface-enhanced Raman Spectroscopy (SERS), and biosensors [50]. Molecular-based methods, including Cluster Amplification Pattern (CAP) and multiplex PCR analysis, provide accurate ways to determine the genetic capability of fungi for toxins production by monitoring deletions in the aflatoxin biosynthesis gene cluster [51].

9. Management of aflatoxin production

Mycotoxin detoxification and prevention are done through a method that encompasses both the post-harvest and pre-harvest phases. The pre-harvest period is characterized by emphasizing proactive control initiatives like the cultivation of resistant crop varieties. Adoption of good farming practices, fertilization systems, and optimization of irrigation, effective pest management, and harvesting at the right time to reduce mycotoxin formation. After harvest, detoxification processes use three methodologies: chemical, where the use of ozone and chitosan is involved; biological, where microbial degradation pathways are exploited; and physical, where controlled drying processes and sorting are used. Other new ways that use applications of nanotechnology, plant-based extracts, and novel binding agents are developing the field and can be integrated into a multi-layered defense mechanism against contamination of mycotoxin in the food production chain [52].

Prevention by Good Agricultural Practices (GAPs), biocontrol (as atoxigenic *Aspergillus* strains), awareness campaigns, and predictive modeling, chemical (ozone and ammonization), bio-based (microbial degradation), and physical (irradiation, sorting) processes are applied to reduce aflatoxin in contaminated products [27]. Chemical (e.g., oxidation), physical (e.g., drying and adsorption), and biological (e.g., transgenic crops, competitive inhibition) are traits of aflatoxin contamination reduction. Protective effects are reported with feed additives and antioxidants (e.g., clay binders and polyphenols) [53]. To address contamination while maintaining food quality and nutritional integrity, researchers have developed three primary categories of mycotoxin reduction strategies [54].

Current management strategies involve a combination of physical, chemical, and biological control methods, with a growing emphasis on eco-friendly alternatives like

biopesticides and natural plant extracts to prevent contamination and mitigate economic losses [55]. Effective management necessitates an integrated "farm-to-fork" approach, utilizing pre-harvest strategies like biological control with atoxigenic strains and crop resistance, alongside post-harvest interventions such as rapid drying, sorting, and innovative enzymatic or chemical detoxification [41].

9.1. Chemical Control of aflatoxin production

Chemical control in the production of aflatoxin has a lot of agents, such as inorganic and organic acids, oxidizing agents, alkalis, synthetic antioxidants, and fungicides [56]. One of the most effective and cost-effective methods of decontaminating feed is ammonium solutions or ammonification with gaseous ammonia, which irreversibly changes the molecular structure of the toxin [41]. The alkaline treatment process of nixtamalization using calcium hydroxide is an effective way of reducing the concentration of aflatoxin in maize by opening the lactone rings, in conditions of acidity such as digestion [57, 58].

Organic acids like propionic acids, tartaric, lactic, and citric have high inhibition rates (up to 92%) and convert aflatoxin B₁ to less toxic derivatives [48]. The most effective agents that can degrade AFs are oxidizing agents, like ozone, which reacts electrophilically with the C8–C9 double bond of the furan ring, and the reaction is efficient (up to 100 %) when alkaline pH levels are present [6]. The use of fungicides (cyprodinil, azoxystrobin, and fludioxonil) to prevent conidia germination and fungal growth, and a mixture of fludioxonil and cyprodinil is reported to be the most effective in decreasing contamination by up to 83% [38]. Synthetic antioxidants like butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) help in preventing the growth of *A. flavus* [56, 59]. Zeolites activated by charcoal and other types of clays, like hydrated sodium calcium aluminosilicate (HSCAS) and bentonite, are used as chemical adsorbents that are added into the diet of animals to prevent absorption into the blood by binding toxins in the gastrointestinal tract [41, 42].

Antioxidant-based control measures against the production of aflatoxin, mainly based on oxidative stress, are a prerequisite to toxin production and fungal growth [60]. expressive of a defensive response to scavenge reactive oxygen species (ROS) and mitigate cellular damage caused by environmental

Review Article

stressors like drought and heat, aflatoxin biosynthesis is triggered by aflatoxigenic fungi, like *Aspergillus* section *Flavi* [61]. Consequently, the biological need of the fungus to produce toxins is reduced by the application of antioxidant compounds into the crop environment, thereby reducing the oxidative stress [37]. Propyl paraben (PP), BHT, and BHA are synthetic phenolic antioxidants which have great efficacy as fungi-toxicants. PP and BHA are effective in achieving 100 % inhibition of fungal growth as well as aflatoxin B₁ accumulation at concentrations of 10–20 mmol/L [62]. The application of BHT needs caution, since sub-inhibitory levels of the compound (e.g., 100 mg/L) have been observed to promote the aflatoxin production, which could be more than 6 times higher than untreated controls [63]. Antioxidants help in interfering with mitochondrial respiration, changing the cell membrane permeability, and suppressing the expression of the aflatoxin biosynthetic gene cluster, including *aflR* and *aflS* genes [1].

Phenolic acids and natural phytochemicals provide food-grade options for aflatoxin management [12]. Caffeic acid shows a strong anti-aflatoxigenic effect, which is able to decrease production of aflatoxin by over 95% without affecting fungal growth through inducing alkyl hydroperoxide reductases that deactivate organic peroxides [37]. Ferulic acid and trans-cinnamic acid are effective in inhibiting *A. parasiticus* and *A. flavus* populations in maize, and particular combinations show inhibition of aflatoxin B₁ under various water activity levels [12]. The other natural antioxidants, like tea polyphenols, gallic acid in walnut tannins, and isoflavones in peanuts, have also been shown to inhibit the production of toxin by regulating fungal oxidative stress responses [64].

Selenium (Se) biofortification is a new combined approach, which regulates the concentrations of aflatoxin and increases the nutritional value of crops [11]. Se does not inhibit the growth of fungi, but a non-lethal dose (0.86 µg/g) leads to a decrease in the maximum aflatoxin B₁ production by toxigenic strains. Antioxidant treatment by Se has been shown to enhance the Darwinian fitness of atoxigenic strains relative to toxigenic counterparts, which is a critical property to biocontrol strategies of competitive exclusion [11]. Additional layers of control are offered by biological extracts and essential oils (EOs) that have high antioxidant activity [60].

Oil of clove, rich in eugenol (83.25%), and blend essential oils

of oregano, cinnamon, and lemongrass demonstrate synergistic inhibitory effects on toxin production and fungal growth by inducing ultrastructural damage to the hyphae and silencing early biosynthetic genes, such as *aflD* and *aflC* [30]. Lactic acid bacteria (LAB) have a role in bio-protection through the release of a cell-free supernatant (CFS) high in peptides and phenolic acids that supplement the functional antioxidants of crops like maize, which is effective in protecting against fungal infestation [15]. Pok et al. [65] tested the ability of three citrus flavonoids; neohesperidin, naringin, and quercetin to inhibit the accumulation of AF in maize contaminated with *A. parasiticus* and found an effective combination of flavonoids (0.24 mM neohesperidin, 0.39 mM naringin, and 0.40 mM quercetin), when applied to maize at 0.95 water activity (a_w), reduced AFB₁, AFG₁, AFB₂, and AFG₂ accumulation by 85% to 100%. This same mixture also shows high effects at a higher 0.98 a_w , achieving 93% to 98% reduction. Table (2) presents chemical agents as a strategy for reducing AFs in maize.

10. Conclusion

Aflatoxin contamination in maize remains a critical global issue due to its significant health risks and economic impacts, but the use of antioxidants presents a promising multi-faceted strategy for mitigation. Research indicates that oxidative stress acts as a prerequisite for aflatoxin biosynthesis, and therefore, the introduction of antioxidants, including elements like selenium, phenolic acids such as caffeic acid, and synthetic compounds like BHA or propyl paraben, can effectively suppress toxin production by alleviating this stress. Furthermore, certain antioxidants can selectively enhance the fitness of atoxigenic strains over toxigenic counterparts, which facilitate more effective biological control through competitive exclusion. While natural extracts, such as plant-based essential oils and organic acids, offer eco-friendly alternatives to traditional synthetic fungicides, the timing and concentration of their application are vital for efficacy. Ultimately, integrating antioxidant treatments into broader integrated pest management strategies, particularly through pre-harvest biofortification and improved post-harvest storage practices, is essential for ensuring the safety and nutritional quality of maize supplies worldwide.

Review Article

Table 2. The use of chemical control agents for reducing AF contamination by toxigenic fungi in maize substrates across affecting AFs-producing fungus and biosynthesis, and through decontamination of accumulated AFs.

Substrate	Toxigenic fungus	Initial toxin concentration	Chemical agent	Applied concentration	Toxin reduction (%)	Reference
Chemical control of aflatoxigenic fungi and AF biosynthesis						
Maize	<i>Aspergillus flavus</i> RCM89	109.65 ppm AFB ₁	trans-cinnamic acid (CA)	25 mM	100%	[11]
			ferulic acid (FA)	30 mM		
			CA-FA mixture	25 + 30 mM		
Maize kernels	<i>A. flavus</i>	109.65 ppb AFB ₁	trans-cinnamic acid (CA)	25 mM	100%	[12]
			Ferulic acid (FA)	30 mM		
			trans-cinnamic acid (CA) + Ferulic acid (FA)	25 mM + 30 mM		
Maize	<i>A. parasiticus</i>	0.12 ppm AFs	Propionic acid	0.25%	100%	[36]
		1.19 ppm AFs		0.5%		
		0.15 ppm AFs	Benzoic acid	0.3%		
		1.34 ppm AFs		0.5%		
		0.19 ppm AFs	Tartaric acid	0.4%		
Maize kernels	<i>A. flavus</i>	27.5 - 32.3 µg/kg AFB ₁	Citric acid (in biopolymer formulation)	0.5%	81%	[67]
Maize	<i>A. flavus</i> NRRL 3357	1.95 µg/g AFB ₁	Cinnamon, Oregano, Lemongrass Composite Essential Oil (COL-CEO)	0.6 - 1.0 µL/disc	67.53% - 72.68%	[61]
Maize	<i>A. flavus</i>	83 µg/kg AFB ₁	Ozone	90 mg/L for 20-40 min	78% - 88%	[6]
Maize	<i>A. flavus</i>	0.01 µg/L AFB ₁	Curcumin and blue light (430 nm)	1000 µM (for 14 days)	100%	[70]
Chemical control of accumulated AFs						
Maize	<i>A. flavus</i>	1000 ng/mL AFB ₁ and 250 ng/mL AFB ₂	Vanillic acid + laccase	1 mM	26% for AFB ₁ and 26.6% for AFB ₂	[66]
Maize	<i>A. flavus</i> and <i>A. parasiticus</i>	0.01 µg/L AFs	Ammonium bicarbonate	0.80%	100%	[68]
Maize flour	<i>A. flavus</i>	53.60 µg/kg AFB ₁	Ozone (O3)	75 mg/L (for 60 min)	78.76%	
Maize	<i>A. flavus</i>	56.91 - 61.66 µg/kg AFB ₁	Ozone	6 mg/hour	100%	[58]
Maize	<i>A. flavus</i>	358.6 µg/kg AFB ₁	Free radicals (via power ultrasound)	1.65 W/cm ³	21.2%	[69]

*ppm: part per million.

Review Article

Author's contributions

Marwa A. Mahmoud, Abdallah M. A. Hassane, and Nageh F. Abo-Dahab: Conceptualization, literature review, writing-original draft, preparation graphics and software, writing-reviewing and editing.

Conflict of interest statement

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding

No funding was received for writing this manuscript.

Author information

Corresponding authors: Abdallah M. A. Hassane*

E-mail: abdallahhassane@azhar.edu.eg

ORCID iD: [0000-0003-0820-8087](https://orcid.org/0000-0003-0820-8087)

Data Availability

No such data is used.

References

- [1] Yu, Y., Li, F., Cheng, Z., Li, M., Xu, Z., Sun, L., ... & Yang, K. (2025). Synergistic inhibition of *Aspergillus flavus* by organic acid salts: growth, oxidative stress, and aflatoxin gene modulation. *Frontiers in Veterinary Science*, 12, 1608792. <https://doi.org/10.3389/fvets.2025.1608792>
- [2] Noman, S. J. (2022). Mycotoxins in food. *Egyptian Academic Journal of Biological Sciences, G. Microbiology*, 14(2), 19-28. <https://doi.org/10.21608/eajbsg.2022.255915>
- [3] Ahmad, M. M., Qamar, F., Saifi, M., & Abdin, M. Z. (2022). Natural inhibitors: A sustainable way to combat aflatoxins. *Frontiers in Microbiology*, 13, 993834. <https://doi.org/10.3389/fmicb.2022.993834>
- [4] Farghl, A. A. M., El-Sheekh, M. M., & El-Shahir, A. A. (2023). Seaweed extracts as biological control of aflatoxins produced by *Aspergillus parasiticus* and *Aspergillus flavus*. *Egyptian Journal of Biological Pest Control*, 33(1), 50. <https://doi.org/10.1186/s41938-023-00696-x>
- [5] Nsabiymva, G., Mutege, C. K., Wagacha, J. M., Mohamed, A. B., Njeru, N. K., Ndayihanzamaso, P., ... & Bandyopadhyay, R. (2023). Aflatoxin contamination of maize and groundnut in Burundi: Distribution of contamination, identification of causal agents and potential biocontrol genotypes of *Aspergillus flavus*. *Frontiers in microbiology*, 14, 1106543. <https://doi.org/10.3389/fmicb.2023.1106543>
- [6] Akullo, J. O., Okello, D. K., Mohammed, A., Muyinda, R., Amayo, R., Magumba, D., ... & Mweetwa, A. (2025). A comprehensive review of aflatoxin in groundnut and maize products in Africa: Prevalence, detection and mitigation strategies. *Journal of Food Quality*, 1(2025), 2810946. <https://doi.org/10.1155/jfq/2810946>
- [7] Nji, Q. N., Babalola, O. O., & Mwanza, M. (2022). Aflatoxins in maize: Can their occurrence be effectively managed in Africa in the face of climate change and food insecurity?. *Toxins*, 14(8), 574. <https://doi.org/10.3390/toxins14080574>
- [8] Gachara, G., Suleiman, R., Kilima, B., Taoussi, M., El Kadili, S., Fauconnier, M. L., ... & Lahlali, R. (2024). Pre-and post-harvest aflatoxin contamination and management strategies of *Aspergillus* spoilage in East African Community maize: review of etiology and climatic susceptibility. *Mycotoxin Research*, 40(4), 495-517. <https://doi.org/10.1007/s12550-024-00555-0>
- [9] Kelman, M. J., Miller, J. D., Renaud, J. B., Baskova, D., & Sumarah, M. W. (2024). A multi-year study of mycotoxin co-occurrence in wheat and corn grown in Ontario, Canada. *Toxins*, 16(8), 372. <https://doi.org/10.3390/toxins16080372>
- [10] Ostrý, V., Malíř, F., & Pfohl-Leszkowicz, A. (2015). Comparative data concerning aflatoxin contents in Bt maize and non-Bt isogenic maize in relation to human and animal health – a review. *Acta Veterinaria Brno*, 84(1), 47–53. <https://doi.org/10.2754/avb201585010047>
- [11] Katati, B., Kovacs, S., Njapau, H., Kachapulula, P. W., Zwaan, B. J., van Diepeningen, A. D., & Schoustra, S. E. (2023). Aflatoxigenic *Aspergillus* modulates aflatoxin-B1 levels through an antioxidative mechanism. *Journal of Fungi*, 9(6), 690. <https://doi.org/10.3390/jof9060690>
- [12] Nesci, A., Gsponer, N., & Etcheverry, M. (2007). Natural maize phenolic acids for control of aflatoxigenic fungi on maize. *Journal of Food Science*, 72(5), M180-M185. <https://doi.org/10.1111/j.1750-3841.2007.00394.x>
- [13] Aziz, S. Y., & Bean, G. A (1995). Influence of antioxidant and antifungal compounds on toxigenic fungi in

Review Article

- oilseeds. Egyptian Journal of Agricultural Research, 73(3), 795-809. <https://doi.org/10.21608/ejar.1995.442181>
- [14] Saleh, H. M., Shetaia, Y. H., Fahmy, H. A., & Mahdy, H. (2015). Antifungal activity of some lactic acid bacteria against aflatoxin-producing aspergilli. Egyptian Journal of Agricultural Sciences, 66(1), 91-104. <https://doi.org/10.21608/ejarc.2015.213220>
- [15] Salman, M., Javed, M. R., Ali, H., Mustafa, G., Tariq, A., Sahar, T., ... & Tawab, A. (2022). Bioprotection of *Zea mays* L. from aflatoxigenic *Aspergillus flavus* by *Loigolactobacillus coryniformis* BCH-4. *PloS one*, 17(8), e0271269. <https://doi.org/10.1371/journal.pone.0271269>
- [16] Smaoui, S., D'Amore, T., Tarapoulouzi, M., Agriopoulou, S., & Varzakas, T. (2023). Aflatoxins contamination in feed commodities: From occurrence and toxicity to recent advances in analytical methods and detoxification. *Microorganisms*, 11(10), 2614. <https://doi.org/10.3390/microorganisms11102614>
- [17] Kumar, P., Gupta, A., Mahato, D. K., Pandhi, S., Pandey, A. K., Kargwal, R., ... & Harakeh, S. (2022). Aflatoxins in cereals and cereal-based products: Occurrence, toxicity, impact on human health, and their detoxification and management strategies. *Toxins*, 14(10), 687. <https://doi.org/10.3390/toxins14100687>
- [18] Agriopoulou, S., Stamatelopoulou, E., & Varzakas, T. (2020). Advances in occurrence, importance, and mycotoxin control strategies: Prevention and detoxification in foods. *Foods*, 9(2), 137. <https://doi.org/10.3390/foods9020137>
- [19] Dai, C., Tian, E., Hao, Z., Tang, S., Wang, Z., Sharma, G., ... & Shen, J. (2022). Aflatoxin B1 toxicity and protective effects of curcumin: Molecular mechanisms and clinical implications. *Antioxidants*, 11(10), 2031. <https://doi.org/10.3390/antiox11102031>
- [20] Khan, R., Anwar, F., & Ghazali, F. M. (2024). A comprehensive review of mycotoxins: Toxicology, detection, and effective mitigation approaches. *Heliyon*, 10(8), e28361. <https://doi.org/10.1016/j.heliyon.2024.e28361>
- [21] Abrehame, S., Manoj, V. R., Hailu, M., Chen, Y. Y., Lin, Y. C., & Chen, Y. P. (2023). Aflatoxins: Source, detection, clinical features and prevention. *Processes*, 11(1), 204. <https://doi.org/10.3390/pr11010204>
- [22] Okechukwu, V. O., Adelusi, O. A., Kappo, A. P., Njobeh, P. B., & Mamo, M. A. (2024). Aflatoxins: Occurrence, biosynthesis, mechanism of action and effects, conventional/emerging detection techniques. *Food Chemistry*, 436, 137775. <https://doi.org/10.1016/j.foodchem.2023.137775>
- [23] Atehnkeng, J., Ojiambo, P. S., Ortega-Beltran, A., Augusto, J., Cotty, P. J., & Bandyopadhyay, R. (2022). Impact of frequency of application on the long-term efficacy of the biocontrol product Aflasafe in reducing aflatoxin contamination in maize. *Frontiers in Microbiology*, 13, 1049013. <https://doi.org/10.3389/fmicb.2022.1049013>
- [24] Hassane, A. M. A. (2017). Evaluation of advanced analytical methods for detection of aflatoxins and aflatoxigenic fungi in some bakery's flour samples in Upper Egypt [Unpublished Doctoral thesis], Al-Azhar University, Egypt.
- [25] Yu, J., Chang, P. K., Ehrlich, K. C., Cary, J. W., Bhatnagar, D., Cleveland, T. E., ... & Bennett, J. W. (2004). Clustered pathway genes in aflatoxin biosynthesis. *Applied and Environmental Microbiology*, 70(3), 1253-1262. <https://doi.org/10.1128/aem.70.3.1253-1262.2004>
- [26] Caceres, I., Al Khoury, A., El Khoury, R., Lorber, S., P. Oswald, I., El Khoury, A., ... & Bailly, J. D. (2020). Aflatoxin biosynthesis and genetic regulation: A review. *Toxins*, 12(3), 150. <https://doi.org/10.3390/toxins12030150>
- [27] Jallow, A., Xie, H., Tang, X., Qi, Z., & Li, P. (2021). Worldwide aflatoxin contamination of agricultural products and foods: From occurrence to control. *Comprehensive Reviews in Food Science and Food Safety*, 20(3), 2332-2381. <https://doi.org/10.1111/1541-4337.12734>
- [28] Passone, M. A., Resnik, S. L., & Etcheverry, M. G. (2005). In vitro effect of phenolic antioxidants on germination, growth and aflatoxin B1 accumulation by peanut *Aspergillus* section Flavi. *Journal of Applied Microbiology*, 99(3), 682-691. <https://doi.org/10.1111/j.1365-2672.2005.02661.x>
- [29] Saber, S. M., Youssef, M. S., Arafa, R. F. & Hassane, A. M. A. (2016). Mycotoxins production by *Aspergillus ostianus* Wehmer and using phytochemicals as control agent. *Journal of Science and Engineering Research*, 3(2), 198-213.
- [30] Achar, P. N., Quyen, P., Adukwu, E. C., Sharma, A., Msimanga, H. Z., Nagaraja, H., & Sreenivasa, M. Y. (2020). Investigation of the antifungal and anti-aflatoxigenic potential

Review Article

of plant-based essential oils against *Aspergillus flavus* in peanuts. *Journal of Fungi*, 6(4), 383. <https://doi.org/10.3390/jof6040383>

[31] Hussain, A., & Shafqatullah, J. A., Zia-ur-Rehman (2012). 02. Inhibition of aflatoxin producing fungus growth using chemical, herbel compounds/spices and plants. *Pure and Applied Biology (PAB)*, 1(1), 8-13.

[32] Dahab, N. F. A., Abdel-Hadi, A. M., Abdul-Raouf, U. M., El-Shanawany, A. A., & Hassane, A. M. (2016). Qualitative detection of aflatoxins and aflatoxigenic fungi in wheat flour from different regions of Egypt. *IOSR Journal of Environmental Science, Toxicology and Food Technology (IOSR-JESTFT)*, 10(7), 20-26.

[33] Abdel-Nasser, A., Fathy, H. M., Badr, A., Hathout, A., & Barakat, O. S. (2022). Prevalence of aflatoxigenic fungi in cereal grains and their related chemical metabolites. *Egyptian Journal of Chemistry*, 65(10), 455-470. <https://doi.org/10.21608/ejchem.2022.122494.5487>

[34] Hassane, A. M. A., El-Shanawany, A. A., Abo-Dahab, N. F., Abdel-Hadi, A. M., Abdul-Raouf, U. M., & Mwanza, M. (2017). Influence of different moisture contents and temperature on growth and production of aflatoxin B1 by a toxigenic *Aspergillus flavus* isolate in wheat flour. *Journal of ecology of health & environment*, 5(3), 77-83. <https://doi.org/10.18576/jehe/050302>

[35] Passone, M. A., Resnik, S., & Etcheverry, M. G. (2007a). Antiaflatoxigenic property of food grade antioxidants under different conditions of water activity in peanut grains. *International Journal of Food Microbiology*, 118(1), 8–14. <https://doi.org/10.1016/j.ijfoodmicro.2007.05.012>

[36] Singh, R., Mandal, A. B., & Biswas, A. (2017). Efficacy of propionic, benzoic and tartaric acids in preventing biosynthesis of aflatoxins in poultry feed. *Animal Nutrition and Feed Technology*, 17(1), 157-164. <https://doi.org/10.5958/0974-181x.2017.00016.6>

[37] Kim, J. H., Yu, J., Mahoney, N., Chan, K. L., Molyneux, R. J., Varga, J., ... & Campbell, B. C. (2008). Elucidation of the functional genomics of antioxidant-based inhibition of aflatoxin biosynthesis. *International Journal of Food Microbiology*, 122(1–2), 49–60. <https://doi.org/10.1016/j.ijfoodmicro.2007.11.058>

[38] Lagogianni, C. S., & Tsitsigiannis, D. I. (2018). Effective chemical management for prevention of aflatoxins in maize.

Phytopathologia Mediterranea, 57(1), 186-198. https://doi.org/10.14601/Phytopathol_Mediterr-22492

[39] Moon, Y.-S., Kim, H.-M., Chun, H. S., & Lee, S.E., (2018). Organic acids suppress aflatoxin production via lowering expression of aflatoxin biosynthesis-related genes in *Aspergillus flavus*. *Food Control*, 88, 207–216. <https://doi.org/10.1016/j.foodcont.2018.01.017>

[40] Fusilier, K., Chilvers, M. I., Limay-Rios, V., & Singh, M. P. (2022). Mycotoxin co-occurrence in Michigan harvested maize grain. *Toxins*, 14(7), 431. <https://doi.org/10.3390/toxins14070431>

[41] Gemedé, H. F. (2025). Toxicity, mitigation, and chemical Analysis of aflatoxins and other toxic metabolites produced by *Aspergillus*: A comprehensive review. *Toxins*, 17(7), 331. <https://doi.org/10.3390/toxins17070331>

[42] Jiang, Y., Ogunade, I. M., Vyas, D., & Adesogan, A. T. (2021). Aflatoxin in dairy cows: toxicity, occurrence in feedstuffs and milk and dietary mitigation strategies. *Toxins*, 13(4), 283. <https://doi.org/10.3390/toxins13040283>

[43] Gnonlonfin, B. G. J., Traore, A. S., & Samake, S. (2019). Effective Control of aflatoxin contamination in staple maize food crop in sub-Saharan Africa: A review of current pre- and postharvest low-cost technologies and perspectives. *Journal of Agricultural Science and Technology*, 9(3), 135-151. <https://doi.org/10.17265/2161-6264/2019.03.001>

[44] Falade, T. D. O., Neya, A., Bonkougou, S., Dagno, K., Basso, A., Senghor, A. L., ... & Bandyopadhyay, R. (2022). Aflatoxin contamination of maize, groundnut, and sorghum grown in Burkina Faso, Mali, and Niger and Aflatoxin exposure assessment. *Toxins*, 14(10), 700. <https://doi.org/10.3390/toxins14100700>

[45] Shekhar, M., Singh, N., Dutta, R., Kumar, S., & Mahajan, V. (2017). Comparative study of qualitative and quantitative methods to determine toxicity level of *Aspergillus flavus* isolates in maize. *PLoS ONE*, 12(12), e0189760. <https://doi.org/10.1371/journal.pone.0189760>

[46] Hassane, A. M. A., El-Shanawany, A. A., Abo-Dahab, N. F., Abdel-Hadi, A. M., Abdul-Raouf, U. M. & Mwanza, M. (2017). Cultural and analytical assays for aflatoxin B production by *Aspergillus flavus* isolates. *Journal of Natural Product Chemistry*, 1(1), 17-23. <https://api.semanticscholar.org/CorpusID:103534353>

Review Article

- [47] Han, S., Yang, Y., Chen, T., Yang, B., Ding, M., Wen, H., ... & Peng, D. (2024). Quantitative determination of aflatoxin B1 in maize and feed by ELISA and time-resolved fluorescent immunoassay based on monoclonal antibodies. *Foods*, 13(2), 319. <https://doi.org/10.3390/foods13020319>
- [48] Shabeer, S., Asad, S., Jamal, A., & Ali, A. (2022). Aflatoxin contamination, its impact and management strategies: An updated review. *Toxins*, 14(5), 307. <https://doi.org/10.3390/toxins14050307>
- [49] Kizis, D., Vichou, A.-E., & Natskoulis, P. I. (2021). Recent advances in mycotoxin analysis and detection of mycotoxigenic fungi in grapes and derived products. *Sustainability*, 13(5), 2537. <https://doi.org/10.3390/su13052537>
- [50] Gabbitas, A., Ahlborn, G., Allen, K., & Pang, S. (2023). Advancing mycotoxin detection: Multivariate rapid analysis on corn using surface enhanced raman spectroscopy (SERS). *Toxins*, 15(10), 610. <https://doi.org/10.3390/toxins15100610>
- [51] Barošević, T., Bagi, F., Savić, Z., Ljubičić, N., & Ivanović, I. (2022). Assessment of maize hybrids resistance to aspergillus ear rot and aflatoxin production in environmental conditions in Serbia. *Toxins*, 14(12), 887. <https://doi.org/10.3390/toxins14120887>
- [52] Awuchi, C. G., Ondari, E. N., Ogbonna, C. U., Upadhyay, A. K., Baran, K., Okpala, C. O. R., & Guiné, R. P. F. (2021). Mycotoxins affecting animals, foods, humans, and plants: Types, occurrence, toxicities, action mechanisms, prevention, and detoxification strategies—A review. *Foods*, 10(6), 1279. <https://doi.org/10.3390/foods10061279>
- [53] Popescu, R. G., Rădulescu, A. L., Georgescu, S. E., & Dinischiotu, A. (2022). Aflatoxins in feed: Types, metabolism, health consequences in swine and mitigation strategies. *Toxins*, 14(12), 853. <https://doi.org/10.3390/toxins14120853>
- [54] Tiwari, S., Shrestha, K., Dhimal, M., Timsina, J., Belbase, K., & Dahal, P. (2022). Complementing food and nutrition security using toxin minimizing dry chain and integrated pest management: A review. *Nepal Journal of Science and Technology*, 20(2), 131–146. <https://doi.org/10.3126/njst.v20i2.45804>
- [55] Abdel-Razek, A. G., Badr, A. N., & Shehata, M. G. (2017). Characterization of olive oil by-products: Antioxidant activity, its ability to reduce aflatoxigenic fungi hazard and its aflatoxins. *Annual Research & Review in Biology*, 14(5), 1–14. <https://doi.org/10.9734/ARRB/2017/35065>
- [56] Konlan, S. Y., Konyannik, B. Y., Benson-Obeng, G., Koomson, M., Akomanyi, B. N., Bamfo, B. S., ... & Osei, G. (2025). Management practices to mitigate aflatoxin contamination in maize: A review. *Asian Research Journal of Agriculture*, 18(3), 461–476. <https://doi.org/10.9734/arja/2025/v18i3755>
- [57] Augusto, J., Atehnkeng, J., Ortega-Beltran, A., Cotty, P. J., & Bandyopadhyay, R. (2024). Keeping toxigenic *Aspergillus* section Flavi and aflatoxin contamination at bay by deploying atoxigenic-based biocontrol products during production of groundnut and maize in Mozambique. *Frontiers in Microbiology*, 15, 1501924. <https://doi.org/10.3389/fmicb.2024.1501924>
- [58] Kinyoro, I. S., & Kaale, L. D. (2025). Aflatoxin decontamination of artificially contaminated maize by ozone and microwave at different PH. *Applied Food Research*, 5(1), 100873. <https://doi.org/10.1016/j.afres.2025.100873>
- [59] Hassane A. M. A., Abo-Dahab N. F., El-Shanawany A. A., Abdel-Hadi A. M., Abdul-Raouf U. M., Ali A. M. & Sultan Y. Y. (2018). *In vitro* and *in situ* impact of safe synthetic and natural antioxidants on populations and aflatoxin B₁ accumulation by *Aspergillus flavus*. *Journal of Biotechnology Science Research*, 5(1), 1-16.
- [60] El Khoury, R., Atoui, A., Mathieu, F., Kawtharani, H., El Khoury, A., Maroun, R. G., & El Khoury, A. (2017). Antifungal and antiochratoxigenic activities of essential oils and total phenolic extracts: A comparative study. *Antioxidants*, 6(3), 44. <https://doi.org/10.3390/antiox6030044>
- [61] Xiang, F., Zhao, Q., Zhao, K., Pei, H., & Tao, F. (2020). The Efficacy of composite essential oils against aflatoxigenic fungus *Aspergillus flavus* in maize. *Toxins*, 12(9), 562. <https://doi.org/10.3390/toxins12090562>
- [62] Passone, M. A., Resnik, S., & Etcheverry, M. G. (2007). Potential use of phenolic antioxidants on peanut to control growth and aflatoxin B1 accumulation by *Aspergillus flavus* and *Aspergillus parasiticus*. *Journal of the Science of Food and Agriculture*, 87(11), 2121–2130. <https://doi.org/10.1002/jsfa.2975>
- [63] Pupovac-Velikonja, A., Velikonja, J., & Drrigl, A. (1986). Kultur von *Aspergillus parasiticus* im Apfelsaft.

Review Article

Zeitschrift für Lebensmittel-Untersuchung und Forschung, 182(4), 303–306. <https://doi.org/10.1007/BF01027629>

[64] Mohsen, E., El-Metwally, M. A., Ibrahim, A. A., & Soliman, M. I. (2023). Impact of green antioxidants on decreasing the aflatoxins percentage in peanut oil seed (*Arachis hypogaea* L.) during storage. Science Progress, 106(2), 1-18. <https://doi.org/10.1177/00368504231176165>

[65] Pok, P. S., García Londoño, V. A., Vicente, S., Romero, S. M., Pacín, A., Tolaba, M., ... & Resnik, S. L. (2020). Evaluation of citrus flavonoids against *Aspergillus parasiticus* in maize: Aflatoxins reduction and ultrastructure alterations. Food Chemistry, 318, 126414. <https://doi.org/10.1016/j.foodchem.2020.126414>

[66] Bossa, M., Alaniz-Zanon, M. S., Monesterolo, N. E., Monge, M. D. P., Coria, Y. M., Chulze, S. N., & Chiotta, M. L. (2024). Aflatoxin decontamination in maize steep liquor obtained from bioethanol production using laccases from species within the basidiomycota phylum. Toxins, 16(1), 27. <https://doi.org/10.3390/toxins16010027>

[67] Rhein, P., Bossa, M., Monge, M. D. P., Giovanini, D., Barbero, C. A., Chulze, S. N., ... & Alaniz-Zanon, M. S. (2026). Reducing aflatoxin accumulation in maize: Development and performance of a novel biological input. Toxins, 18(1), 49. <https://doi.org/10.3390/toxins18010049>

[68] Gómez-Salazar, J. A., Ruiz-Hernández, K., Martínez-Miranda, M. M., & Castro-Ríos, K. (2023). Postharvest strategies for decontamination of aflatoxins in cereals. Food Reviews International, 39(7), 3635–3662. <https://doi.org/10.1080/87559129.2021.2013254>

[69] Liu, Y., Liu, Y., Zhao, W., Li, M., Liu, N., & Bian, K. (2022). Reduction of aflatoxin B1 and zearalenone contents in corn using power ultrasound and its effects on corn quality. Toxins, 14(12), 834. <https://doi.org/10.3390/toxins14120834>

[70] Nguenha, R., Damyeh, M. S., Phan, A. D., Hong, H. T., Chaliha, M., O'Hare, T. J., ... & Sultanbawa, Y. (2021). Effect of photosensitization mediated by curcumin on carotenoid and aflatoxin content in different maize varieties. Applied Sciences, 11(13), 5902. <https://doi.org/10.3390/app11135902>